

REVISION: DRUG ABUSE

Drag and drop the correct answers in the box provided below.

Withdrawal symptoms:

1. Restlessness.
2. Excessive sweating.
3. Uncontrolled shaking.
4. Anxiety.
5. Fever.
6. Headaches.

1. Miscarriage in pregnant women.

1. Miscarriage in pregnant women.
2. Lung cancer.
3. Cancer of the mouth, throat, pancreas, kidneys & urinary bladder.
4. Chronic bronchitis.
5. Emphysema.

1. Used to increase muscle tissue.

1. Used to increase muscle tissue.
2. Can be abused as a sports drug.
3. Causes aggression and male infertility.

1. Dry eyes & mouth.
2. Increased hunger.
3. Hallucinations.
4. Appear sleepy.

1. Slurred speech.
2. Blurred vision.
3. Clumsiness.
4. Inability to walk steadily.

1. Increased breathing & heart rates.
2. Dilated pupils.
3. Loss of appetite.

1. Tiredness.
2. Loss of appetite.
3. Very quick addiction to the drug.

Causes a temporary loss of sensation.

To relieve pain without causing any numbness.

Contain tar, nicotine, carbon monoxide & irritants.

For treatment of diseases caused by bacteria.

1. Liver damage.
2. Violence in the family.
3. Crime.

