

ONE MARK TEST

P. ELANGO VAN, B.T. Assistant (Mathematics)
GOVERNMENT HIGHER SECONDARY SCHOOL
KOLIYANUR – VILLUPURAM DISTRICT



ENGLISH MEDIUM

LESSON - 1

TEST - 1

- 1 The range of the relation $R = \{(x, x^2) \mid x \text{ is a prime number less than } 13\}$ is
(A) $\{2, 3, 5, 7\}$ (B) $\{2, 3, 5, 7, 11\}$
(C) $\{4, 9, 25, 49, 121\}$ (D) $\{1, 4, 9, 25, 49, 121\}$
- 2 If $\{(a, 8), (6, b)\}$ represents an identity function, then the value of a and b are respectively
(A) (8, 6) (B) (8, 8) (C) (6, 8) (D) (6, 6)
- 3 If $A = \{1, 2\}$, $B = \{1, 2, 3, 4\}$, $C = \{5, 6\}$ and $D = \{5, 6, 7, 8\}$ then state which of the following statement is true.
(A) $(A \times C) \subset (B \times D)$ (B) $(B \times D) \subset (A \times C)$
(C) $(A \times B) \subset (A \times D)$ (D) $(D \times A) \subset (B \times A)$
- 4 Let f and g be two functions given by
 $f = \{(0, 1), (2, 0), (3, -4), (4, 2), (5, 7)\}$
 $g = \{(0, 2), (1, 0), (2, 4), (-4, 2), (7, 0)\}$ then the range of $f \circ g$ is
(A) $\{0, 2, 3, 4, 5\}$ (B) $\{-4, 1, 0, 2, 7\}$ (C) $\{1, 2, 3, 4, 5\}$ (D) $\{0, 1, 2\}$
- 5 If there are 1024 relations from a set $A = \{1, 2, 3, 4, 5\}$ to a set B , then the number of elements in B is
(A) 3 (B) 2 (C) 4 (D) 8

- 6 Let $A = \{1, 2, 3, 4\}$ and $B = \{4, 8, 9, 10\}$. A function $f : A \rightarrow B$ given by $f = \{(1, 4), (2, 8), (3, 9), (4, 10)\}$ is a
- (A) Many-one function (B) Identity function
(C) One-to-one function (D) Into function
- 7 If $f : A \rightarrow B$ is a bijective function and if $n(B) = 7$, then $n(A)$ is equal to
- (A) 7 (B) 49 (C) 1 (D) 14
- 8 $A = \{a, b, p\}$, $B = \{2, 3\}$, $C = \{p, q, r, s\}$ then $n[(A \cup C) \times B]$ is
- (A) 8 (B) 20 (C) 12 (D) 16
- 9 Let $f(x) = \sqrt{1 + x^2}$ then
- (A) $f(xy) = f(x) \cdot f(y)$ (B) $f(xy) \geq f(x) \cdot f(y)$
(C) $f(xy) \leq f(x) \cdot f(y)$ (D) None of these
- 10 If $n(A \times B) = 6$ and $A = \{1, 3\}$ then $n(B)$ is
- (A) 1 (B) 2 (C) 3 (D) 6