

Name: _____

Class: _____

REVISION: DRUG ABUSE

Drag and drop the correct answers in the box provided below.

Withdrawal symptoms:

1. Restlessness.
2. Excessive sweating.
3. Uncontrolled shaking.
4. Anxiety.
5. Fever.
6. Headaches.

1. Miscarriage in pregnant women.
2. Lung cancer.
3. Cancer of the mouth, throat, pancreas, kidneys & urinary bladder.
4. Chronic bronchitis.
5. Emphysema.

1. Used to increase muscle tissue.
2. Can be abused as a sports drug.
3. Causes aggression and male infertility.

1. Dry eyes & mouth.
2. Increased hunger.
3. Hallucinations.
4. Appear sleepy.

1. Slurred speech.
2. Blurred vision.
3. Clumsiness.
4. Inability to walk steadily.

1. Increased breathing & heart rates.
2. Dilated pupils.
3. Loss of appetite.

1. Tiredness.
2. Loss of appetite.
3. Very quick addiction to the drug.

Causes a temporary loss of sensation.

To relieve pain without causing any numbness.

Contain tar, nicotine, carbon monoxide & irritants.

For treatment of diseases caused by bacteria.

1. Liver damage.
2. Violence in the family.
3. Crime.

