

# Factor

Name \_\_\_\_\_

Date \_\_\_\_\_

The example shows a way of finding common multiples by writing the first ten multiples for each pair of numbers, then circling the common multiples.

| 4  | 6  |
|----|----|
| 4  | 6  |
| 8  | 12 |
| 12 | 18 |
| 16 | 24 |
| 20 | 30 |
| 24 | 36 |
| 28 | 42 |
| 32 | 48 |
| 36 | 54 |
| 40 | 60 |

and you write :

same multiple : 12, 24. and 36

| 2 | 5 |
|---|---|
|   |   |
|   |   |
|   |   |
|   |   |
|   |   |
|   |   |
|   |   |
|   |   |
|   |   |

same multiple is :

| 3 | 4 |
|---|---|
|   |   |
|   |   |
|   |   |
|   |   |
|   |   |
|   |   |
|   |   |
|   |   |
|   |   |

same multiple is :

| 4 | 7 |
|---|---|
|   |   |
|   |   |
|   |   |
|   |   |
|   |   |
|   |   |
|   |   |
|   |   |
|   |   |

same multiple is :

| 3 | 10 |
|---|----|
|   |    |
|   |    |
|   |    |
|   |    |
|   |    |
|   |    |
|   |    |
|   |    |
|   |    |

same multiple is :