

What Price Fresh Flowers?



Flowers have long been symbols of love and caring. People send them to express sympathy, to apologise, or just wish someone well. But today, floriculture - the growing and selling of flowers - is very big business, worth £2.2 billion a year in the UK alone. The majority of the cut-flowers sold there are imported, these days mostly from countries such as Colombia and Kenya. The Netherlands is the traditional centre of flower production in Europe, and remains a major supplier of flowers. In recent years, however, as labour and production costs have soared, attention there has shifted from flower production to flower trading. Meanwhile, the Kenyan cut-flower industry has grown rapidly, and now provides a vital income for around two million people. It is the country's largest agricultural foreign exchange earner after tea, producing £165 million annually.

For the environmentally conscious, it might seem wasteful that a commodity such as flowers should travel halfway around the world before arriving at a supermarket or florist shop. Just as some environmentalists say that it's better to buy fruit and vegetables grown locally, some also advocate the buying of locally-grown flowers. Thanks to globalisation, however, the UK cut-flower industry now supplies just ten per cent of the country's needs. Twenty years ago it was more like half. What's more, it is suggested that reversing this trend would actually have serious environmental consequences.

Research published in 2006 by Cranfield University in the UK showed that the production of Kenyan flowers, including delivery by air freight and truck, resulted in a carbon footprint nearly six times smaller than that caused by the production of Dutch flowers. Kenya has optimal growing conditions and the warm African sun provides heat and light, whereas growers in the Netherlands and other developed countries require significant inputs of gas and electricity to grow flowers year-round in artificial climate-controlled environments. But there are other environmental factors to consider. A vast range of pesticides, fertilisers and fumigants are used in producing cut flowers. Lake Naivasha, the centre of Kenya's flower industry, is the ideal place to grow roses, thanks to its high altitude and abundant sunlight and water. However, environmental damage has resulted from the development that has followed in the wake of floriculture.



Lake Naivasha itself has shrunk to half its original size, with water levels dropping by three metres, fish catches falling and the native hippopotamus feeling the effects of pollution. Meanwhile, transporting flowers over long distances poses its own set of challenges. Roses, for example, have to be shipped by air rather than sea because they require constant refrigeration and wilt quickly. Transporting other types of flowers by sea can also be tricky compared to air freight. Demand is difficult to predict, which means entire shipping containers can seldom be filled with a single species, but mixing flowers is often inadvisable because some varieties emit gases that spoil others. One strategy is to opt for heartier breeds such as carnations and lilies which are easier to ship and require less refrigeration than roses.

Questions 1-8

Do the following statements agree with the information given in the Reading Passage?

Next to each question, 1-8, write

TRUE if the statement agrees with the information

FALSE if the statement contradicts the information

NOT GIVEN if there is no information on this in the passage

- 1 In recent years, cut flowers have become more expensive to grow in the Netherlands.
- 2 More people are employed in the cut-flower industry in Kenya than in Europe.
- 3 Flowers represent Kenya's most valuable agricultural export.
- 4 The UK has seen a marked decline in the proportion of locally-grown cut flowers on sale.
- 5 The Cranfield study concentrated on the environmental effects of transporting cut flowers.
- 6 The Lake Naivasha region produces a range of cut flowers including roses.
- 7 Supplies of some local food items have been affected by the impact of floriculture around Lake Naivasha.
- 8 Transporting cut flowers by sea is generally more successful than using other means of transport.

The effects of light on plant and animal species




Light is important to organisms for two different reasons. Firstly, it is used as a cue for the timing of daily and seasonal rhythms in both plant and animals, and secondly, it is used to assist growth in plants.

Breeding in most organisms occurs during a part of the year only, and so a reliable cue is needed to trigger breeding behaviour. Day length is an excellent cue because it provides a perfectly predictable pattern of change within the year. In the temperate zone in spring, temperatures fluctuate greatly from day to day, but day length increases steadily by a predictable amount. The seasonal impact of day length on physiological responses is called photoperiodism, and the amount of experimental evidence for this phenomenon is considerable. For example, some species of birds' breeding can be induced even in midwinter simply by increasing day length artificially (Wolfson 1964). Other examples of photoperiodism occur in plants. A short day plant flowers when the day is less than a certain critical length. A long day plant flowers after a certain critical day length is exceeded. In both cases, the critical day length differs from species to species. Plants which flower after a period of vegetative growth, regardless of photoperiod, are known as day-neutral plants.

Breeding seasons in animals such as birds have evolved to occupy the part of the year in which offspring have the greatest chances of survival. Before the breeding season begins, food reserves must be built up to support the energy cost of reproduction, and to provide for young birds both when they are in the nest and after fledging. Thus many temperate-zone birds use the increasing day lengths in spring as a cue to begin the nesting cycle, because this is a point when adequate food resources will be assured.

The adaptive significance of photoperiodism in plants is also clear. Short-day plants that flower in spring in the temperate zone are adapted to maximising seedling growth during the growing season. Long-day plants are adapted for situations that require fertilization by insects, or a long period of seed ripening. Short-day plant that flower in the autumn in the temperate zone are able to build up food reserves over the growing season and over winter as seeds. Day-neutral plants have an evolutionary advantage when the connection between the favourable period for reproduction and day length is much less certain. For example, desert annuals germinate, flower and seed whenever suitable rainfall occurs, regardless of the day length.

The breeding season of some plants can be delayed to extraordinary lengths. Bamboos are perennial grasses that remain in a vegetative state for many years and then suddenly flower, fruit and die (Evans 1976). Every bamboo of the species *Chusquea abietifolio* on the island of Jamaica flowered, set seed and died during 1884. The next generation of bamboo flowered and died between 1916 and 1918, which suggests a vegetative cycle of about 31 years. The climatic trigger for this flowering cycle is not-yet-known, but the adaptive significance is clear. The simultaneous production of masses of bamboo seeds (in some cases lying 12 to 15 centimetres deep on the ground) is more than all the seed-eating animals can cope with at the time, so that some seeds escape being eaten and grow up to form the next generation (Evans 1976).



The second reason light is important to organisms is that it is essential for photosynthesis. This is the process by which plants use energy from the sun to convert carbon from soil or water into organic material for growth. The rate of photosynthesis in a plant can be measured by calculating the rate of its uptake of carbon. There is a wide range of photosynthetic responses of plants to variations in light intensity. Some plants reach maximal photosynthesis at one-quarter full sunlight, and others, like sugarcane, never reach a maximum, but continue to increase photosynthesis rate as the light intensity rises.



Plants, in general, can be divided into two groups: shade-tolerant species and shade-intolerant species. This classification is commonly used in forestry and horticulture. Shade-tolerant plants have lower photosynthetic rates and hence have lower growth rates than those of shade-intolerant species. Plant species become adapted to living in a certain kind of habitat, and in the process evolve a series of characteristics that prevent them from occupying other habitats. Grime (1966) suggests that light may be one of the major components directing these adaptations. For example, eastern hemlock seedlings are shade-tolerant. They can survive in the forest understorey under very low light levels because they have a low photosynthetic rate.

Questions 27-33

Do the following statements agree with the information given in Reading Passage **84**?

TRUE if the statement agrees with the information

FALSE if the statement contradicts the information

NOT GIVEN if there is no information on this

27. There is plenty of scientific evidence to support photoperiodism.
28. Some types of bird can be encouraged to breed out of season.
29. Photoperiodism is restricted to certain geographic areas.
30. Desert annuals are examples of long-day plants.
31. Bamboos flower several times during their life cycle.
32. Scientists have yet to determine the cue for Chusquea abietifolia's seasonal rhythm.
33. Eastern hemlock is a fast-growing plant.

Questions 34-40

Complete the sentences.

Choose **NO MORE THAN THREE WORDS** from the passage for each answer.

34. Day length is a useful cue for breeding in areas where are unpredictable.
35. Plants which do not respond to light levels are referred to as
36. Birds in temperate climates associate longer days with nesting and the availability of
37. Plants that flower when days are long often depend on to help them reproduce.
38. Desert annuals respond to as a signal for reproduction.
39. There is no limit to the photosynthetic rate in plants such as
40. Tolerance to shade is one criterion for the of plants in forestry and horticulture.

