















Group A : Match the instructions with the pictures.


Step	Picture	Instruction	
1		a	Pour the buffer into the tube.
2		b	Remove the swab.
3		c	Tilt your child's head back 70 degrees.
4		d	Place the tube in the stand.
5		e	Insert the swab 1.5-2 centimetres into one nostril.
6		f	Open the buffer liquid package.

Group B: Match the instructions with the pictures.

7		g	Move the swab using a circular motion inside the nostril for at least 5 seconds.
8		h	Press on the side of the nose. Rotate the swab for at least 5 seconds.
9		i	Swirl the swab around in the tube
		j	Use the same swab on the other nostril. Follow steps 6-8 on the other nostril.
10		k	Squeeze the swab by pinching the sides of the tube together 10-15 times.
11		l	Insert the swab into the tube.
12			

Group C: Match the instructions with the pictures.

13		m	Turn the tube upside down.
14		n	Put the nozzle on the tube.
15		o	Squeeze 3 drops of the liquid onto the circle on the top of the test.
16		p	Set a timer for 15 minutes.
17		q	Squeeze the tube to get the liquid out of the swab and throw out the swab.
18		r	View the test results after 15 minutes.
19		s	Put the tube back in the stand and wait for 2 minutes.



Information Taken from

https://docs.google.com/presentation/d/1zVDiI-rZ-k9wH-nE34ZelwgtuRjQc3iAOaEjqKOQLg/edit#slide=id.g1031d70bbb9_0_9