

**Example 1.11**

Let  $A = \{1, 2, 3, 4\}$  and  $B = \{2, 5, 8, 11, 14\}$  be two sets. Let  $f: A \rightarrow B$  be a function given by  $f(x) = 3x - 1$ . Represent this function

- (i) by arrow diagram
- (ii) in a table form
- (iii) as a set of ordered pairs
- (iv) in a graphical form

**Solution :**

$$A = \{1, 2, 3, 4\}; B = \{2, 5, 8, 11, 14\}$$

$$f(x) = 3x - 1$$

$$f(1) = 3 \underline{\quad} - 1 = \underline{\quad} - 1 = 2$$

$$f(2) = 3 \underline{\quad} - 1 = \underline{\quad} - 1 = 5$$

$$f(3) = 3 \underline{\quad} - 1 = \underline{\quad} - 1 = 8$$

$$f(4) = 4 \underline{\quad} - 1 = \underline{\quad} - 1 = 11$$

**(i) Arrow diagram**

Let us represent the function  $f: A \rightarrow B$  by



an arrow diagram

**(iii) Set of ordered pairs**

The function  $f$  can be represented as a set of ordered pair as

**(iv) Graphical form**

In the adjacent  $xy$ -plane the points  $(1, 2), (2, 5), (3, 8), (4, 11)$  are plotted



$$f = \{(1, 2), (2, 5), (3, 8), (4, 11)\}$$

**(ii) Table form**

The given function  $f$  can be represented in a tabular form as given below

$x$	1	2	3	4
$f(x)$	2	5	8	11



Edit with WPS Office

2. Let  $f: A \rightarrow B$  be a function define by

$$f(x) = \frac{x}{2} - 1, \text{ where } A = \{2, 4, 6, 10, 12\},$$

$B = \{0, 1, 2, 4, 5, 9\}$ . Represent  $f$  by

- (i) set of ordered pairs ; (ii) a table ;
- (iii) an arrow diagram ; (iv) a graph

*Solution :*

Given

$$f(x) = \frac{x}{2} - 1$$

$$x = 2 \Rightarrow f(\underline{\hspace{1cm}}) = \frac{\underline{\hspace{1cm}}}{2} - 1 = \underline{\hspace{1cm}} - 1 = 0$$

$$x = 4 \Rightarrow f(\underline{\hspace{1cm}}) = \frac{\underline{\hspace{1cm}}}{2} - 1 = \underline{\hspace{1cm}} - 1 = 1$$

$$x = 6 \Rightarrow f(\underline{\hspace{1cm}}) = \frac{\underline{\hspace{1cm}}}{2} - 1 = \underline{\hspace{1cm}} - 1 = 2$$

$$x = 10 \Rightarrow f(\underline{\hspace{1cm}}) = \frac{\underline{\hspace{1cm}}}{2} - 1 = \underline{\hspace{1cm}} - 1 = 4$$

$$x = 12 \Rightarrow f(\underline{\hspace{1cm}}) = \frac{\underline{\hspace{1cm}}}{2} - 1 = \underline{\hspace{1cm}} - 1 = 5$$

(i) Set of order pairs :



(ii) Table :

(iii) Arrow diagram :

(iv) Graph

$$f = \{(2, 0), (4, 1), (6, 2), (10, 4), (12, 5)\}$$

$x$	2	4	6	10	12
$f(x)$	0	1	2	4	5



3. Represent the function  $f = \{(1, 2), (2, 2), (3, 2), (4, 3), (5, 4)\}$  through (i) an arrow diagram (ii) a

table form (iii) a graph

*Solution :*

(i) Arrow Diagram :



(ii) Table Form :

(iii) Graph :



Edit with WPS Office

$X$	1	2	3	4	5
$f(x)$	2	2	2	3	4



Edit with WPS Office