

FORMULA GENERAL

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

RESUELVE

$$25x^2 + 100x + 100 = 0$$

$$a = \boxed{}$$

$$b = \boxed{}$$

$$c = \boxed{}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$x = \frac{-(\boxed{}) \pm \sqrt{(\boxed{})^2 - \boxed{}(\boxed{})(\boxed{})}}{\boxed{}(\boxed{})}$$

$$x = \frac{-\boxed{} \pm \sqrt{\boxed{}\boxed{}\boxed{}}}{\boxed{}}$$

$$x = \frac{-\boxed{} \pm \sqrt{\boxed{}}}{\boxed{}}$$

$$x = \frac{-\boxed{} \pm \boxed{}}{\boxed{}}$$

$$x_1 = \frac{-\boxed{} + \boxed{}}{\boxed{}} = \boxed{}$$

$$x_2 = \frac{-\boxed{} - \boxed{}}{\boxed{}} = \boxed{}$$

RESUELVE

$$4x^2 + 24x + 32 = 0$$

$$a = \boxed{}$$

$$b = \boxed{}$$

$$c = \boxed{}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$x = \frac{-(\boxed{}) \pm \sqrt{(\boxed{})^2 - \boxed{}(\boxed{})(\boxed{})}}{\boxed{}(\boxed{})}$$

$$x = \frac{-\boxed{} \pm \sqrt{\boxed{} \boxed{} \boxed{}}}{\boxed{}}$$

$$x = \frac{-\boxed{} \pm \sqrt{\boxed{}}}{\boxed{}}$$

$$x = \frac{-\boxed{} \pm \boxed{}}{\boxed{}}$$

$$x_1 = \frac{-\boxed{} + \boxed{}}{\boxed{}} = \boxed{}$$

$$x_2 = \frac{-\boxed{} - \boxed{}}{\boxed{}} = \boxed{}$$

RESUELVE

$$2x^2 + 20x + 50 = 0$$

$$a = \boxed{}$$

$$b = \boxed{}$$

$$c = \boxed{}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$x = \frac{-(\boxed{}) \pm \sqrt{(\boxed{})^2 - \boxed{}(\boxed{})(\boxed{})}}{\boxed{}(\boxed{})}$$

$$x = \frac{-\boxed{} \pm \sqrt{\boxed{} \boxed{} \boxed{}}}{\boxed{}}$$

$$x = \frac{-\boxed{} \pm \sqrt{\boxed{}}}{\boxed{}}$$

$$x = \frac{-\boxed{} \pm \boxed{}}{\boxed{}}$$

$$x_1 = \frac{-\boxed{} + \boxed{}}{\boxed{}} = \boxed{}$$

$$x_2 = \frac{-\boxed{} - \boxed{}}{\boxed{}} = \boxed{}$$

RESUELVE

$$2x^2 + 14x + 24 = 0$$

$$a = \boxed{}$$

$$b = \boxed{}$$

$$c = \boxed{}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$x = \frac{-(\boxed{}) \pm \sqrt{(\boxed{})^2 - \boxed{}(\boxed{})(\boxed{})}}{\boxed{}(\boxed{})}$$

$$x = \frac{-\boxed{} \pm \sqrt{\boxed{}\boxed{}\boxed{}}}{\boxed{}}$$

$$x = \frac{-\boxed{} \pm \sqrt{\boxed{}}}{\boxed{}}$$

$$x = \frac{-\boxed{} \pm \boxed{}}{\boxed{}}$$

$$x_1 = \frac{-\boxed{} + \boxed{}}{\boxed{}} = \boxed{}$$

$$x_2 = \frac{-\boxed{} - \boxed{}}{\boxed{}} = \boxed{}$$

RESUELVE

$$4x^2 + 16x + 12 = 0$$

$$a = \boxed{}$$

$$b = \boxed{}$$

$$c = \boxed{}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$x = \frac{-(\boxed{}) \pm \sqrt{(\boxed{})^2 - \boxed{}(\boxed{})(\boxed{})}}{\boxed{}(\boxed{})}$$

$$x = \frac{-\boxed{} \pm \sqrt{\boxed{} \boxed{} \boxed{}}}{\boxed{}}$$

$$x = \frac{-\boxed{} \pm \sqrt{\boxed{}}}{\boxed{}}$$

$$x = \frac{-\boxed{} \pm \boxed{}}{\boxed{}}$$

$$x_1 = \frac{-\boxed{} + \boxed{}}{\boxed{}} = \boxed{}$$

$$x_2 = \frac{-\boxed{} - \boxed{}}{\boxed{}} = \boxed{}$$

RESUELVE

$$5x^2 + 25x + 30 = 0$$

$$a = \boxed{}$$

$$b = \boxed{}$$

$$c = \boxed{}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$x = \frac{-(\boxed{}) \pm \sqrt{(\boxed{})^2 - \boxed{}(\boxed{})(\boxed{})}}{\boxed{}(\boxed{})}$$

$$x = \frac{-\boxed{} \pm \sqrt{\boxed{} \boxed{} \boxed{}}}{\boxed{}}$$

$$x = \frac{-\boxed{} \pm \sqrt{\boxed{}}}{\boxed{}}$$

$$x = \frac{-\boxed{} \pm \boxed{}}{\boxed{}}$$

$$x_1 = \frac{-\boxed{} + \boxed{}}{\boxed{}} = \boxed{}$$

$$x_2 = \frac{-\boxed{} - \boxed{}}{\boxed{}} = \boxed{}$$

RESUELVE

$$6x^2 + 30x + 24 = 0$$

$$a = \boxed{}$$

$$b = \boxed{}$$

$$c = \boxed{}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$x = \frac{-\boxed{} \pm \sqrt{\boxed{}^2 - \boxed{}(\boxed{})(\boxed{})}}{\boxed{}(\boxed{})}$$

$$x = \frac{-\boxed{} \pm \sqrt{\boxed{} \boxed{} \boxed{}}}{\boxed{}}$$

$$x = \frac{-\boxed{} \pm \sqrt{\boxed{}}}{\boxed{}}$$

$$x = \frac{-\boxed{} \pm \boxed{}}{\boxed{}}$$

$$x_1 = \frac{-\boxed{} + \boxed{}}{\boxed{}} = \boxed{}$$

$$x_2 = \frac{-\boxed{} - \boxed{}}{\boxed{}} = \boxed{}$$

RESUELVE

$$x^2 + 6x - 27 = 0$$

$$a = \boxed{}$$

$$b = \boxed{}$$

$$c = \boxed{}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$x = \frac{-(\boxed{}) \pm \sqrt{(\boxed{})^2 - \boxed{}(\boxed{})(\boxed{})}}{\boxed{}(\boxed{})}$$

$$x = \frac{-\boxed{} \pm \sqrt{\boxed{} \boxed{} \boxed{}}}{\boxed{}}$$

$$x = \frac{-\boxed{} \pm \sqrt{\boxed{}}}{\boxed{}}$$

$$x = \frac{-\boxed{} \pm \boxed{}}{\boxed{}}$$

$$x_1 = \frac{-\boxed{} + \boxed{}}{\boxed{}} = \boxed{}$$

$$x_2 = \frac{-\boxed{} - \boxed{}}{\boxed{}} = \boxed{}$$