

VALOR NUMÉRICO

Si $a = 2$ $b = 3$ $c = 4$ $d = \frac{1}{2}$ $m = \frac{2}{3}$ $n = \frac{3}{4}$ $q = \frac{2}{5}$ $t = \frac{1}{4}$ $p = 5$

1) $\frac{a^2 d^2 \sqrt{n+bd}}{ct}$

2) $qt \sqrt[3]{n + \frac{pc+1}{ac}}$

3) $\frac{p^2}{5} + \frac{c^2}{8} - \frac{d^2}{4}$

4) $\frac{d}{n} + \frac{t}{d} - q$

5) $\frac{b^3 m^2}{2\sqrt{n^2 t^2}}$

6) $\frac{3\sqrt[3]{64d^3 c^6}}{2n}$

7) $a = 2$ $b = -1$ $c = -3$
 $\frac{-b + \sqrt{b^2 - 4ac}}{2a}$

8) $a = 3$ $b = 7$ $c = -6$
 $\frac{-b + \sqrt{b^2 - 4ac}}{2a}$

9) $a = 8$ $b = -10$ $c = 2$
 $\frac{-b + \sqrt{b^2 - 4ac}}{2a}$

17) $a = 3$ $b = 4$ $c = \frac{1}{3}$ $d = \frac{1}{2}$ $m = 6$ $n = \frac{1}{4}$
 $\frac{c}{d} - \frac{m}{n} + 2$

Con los mismos valores anteriores ejercicio 17

19) $\sqrt{b} + \sqrt{n} + \sqrt{6m}$

10) $a = 5$ $b = -10$ $c = 5$
 $\frac{-b + \sqrt{b^2 - 4ac}}{2a}$

11) $a = 6$ $b = 8$ $c = -8$
 $\frac{-b + \sqrt{b^2 - 4ac}}{2a}$

12) $a = 5$ $b = 4$ $c = -1$
 $\frac{-b + \sqrt{b^2 - 4ac}}{2a}$

13) $a = -4$ $t = 2$

$\frac{1}{2}at^2 - 4t + 3$

14) $a = -3$ $t = \frac{1}{3}$
 $\frac{1}{2}at^2 - 2t + 2$

15) $a = -\frac{3}{4}$ $t = 4$
 $\frac{1}{2}at^2 + 4t - 5$

Si $a = 2$ $b = 3$ $c = 4$ $d = \frac{1}{2}$ $m = \frac{2}{3}$ $n = \frac{3}{4}$ $q = \frac{2}{5}$ $t = \frac{1}{4}$ $p = 5$

16) $c\sqrt{c} - d\sqrt{16bn} + q\sqrt{apq}$

18) $a = 3$ $b = 4$ $c = \frac{1}{3}$ $d = \frac{1}{2}$ $m = 6$ $n = \frac{1}{4}$
 $\frac{3}{5}c - \frac{1}{2}b + 2d$

valores del ejercicio 17

20) $\frac{12c-a}{2b} - \frac{16n-a}{m} + \frac{1}{d}$