

Read the drugs label below.

GENUINE ASPIRIN
BAYER
ASPIRIN (NSAID) PAIN RELIEVER/FEVER REDUCER

THE WONDER DRUG. 325^{mg}

Safe Pain Relief Plus Lifesaving Benefits COATED TABLETS

Drug Facts

Active ingredient (in each tablet) Aspirin 325 mg (NSAID)
Purposes Pain reliever/fever reducer
Nonsteroidal anti-inflammatory drug

Uses temporarily relieves
● headache ● muscle pain ● toothache
● menstrual pain ● pain and fever of colds ● minor pain of arthritis

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Aspirin may cause a severe allergic reaction which may include:
● hives ● facial swelling ● asthma (wheezing) ● shock
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you
● are age 60 or older
● have had stomach ulcers or bleeding problems
● take a blood thinning (anticoagulant) or steroid drug
● take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
● have 3 or more alcoholic drinks every day while using this product
● take more or for a longer time than directed
Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if
● stomach bleeding warning applies to you
● you have a history of stomach problems, such as heartburn
● you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
● you are taking a diuretic
● you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for
● gout ● diabetes ● arthritis

FOR ADULT USE

- Safety Relieves Everyday Aches & Pains Like Headaches, Muscle Aches, & Body Aches
- Coated for Easier Swallowing
- Can Help Save Your Life During a Heart Attack When Taken as Directed by a Doctor
- Recommended by Doctors for Aspirin Regimen Use

Aspirin is not appropriate for everyone, so be sure to talk to your doctor before you begin an aspirin regimen.

Tamper-Evident: Packets of 2 Tablets, Do not use if packet is open or torn.

50 Packets/100 Tablets

Drug Facts (continued)

Stop use and ask a doctor if
● an allergic reaction occurs. Seek medical help right away.
● you experience any of the following signs of stomach bleeding:
● feel faint ● vomit blood ● have bloody or black stools
● have stomach pain that does not get better
● pain gets worse or lasts more than 10 days
● redness or swelling is present
● fever lasts more than 3 days
● new symptoms occur
● ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
● drink a full glass of water with each dose
● adults and children 12 years and over: take 1 to 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours
● children under 12 years: consult a doctor

Other information
● save carton for full directions and warnings
● store at room temperature

Inactive ingredients carnauba wax*, corn starch, hypromellose, powdered cellulose, triacetin
*may contain this ingredient

Questions or comments? 1-800-331-4536
(Mon - Fri 9AM - 5PM EST) or www.bayeraspirin.com

A. State true or false

1. The active ingredient in Bayer Aspirin is 325 mg of aspirin. (_ _ _ _)
2. The medicine can be used to relieve headache, muscle pain, and toothache. (_ _ _ _)
3. The label says it is safe for children who have flu symptoms. (_ _ _ _)
4. Bayer Aspirin may cause severe allergic reactions such as asthma or shock. (_ _ _ _)
5. Adults are instructed to take 4 tablets every hour. (_ _ _ _)
6. The product should be stored in a refrigerator. (_ _ _ _)
7. Inactive ingredients include carnauba wax and corn starch. (_ _ _ _)
8. Pregnant or breastfeeding women shouldn't ask a health professional before use. (_ _ _ _)
9. Adults can take more than 12 tablets in 24 hours. (_ _ _ _)
10. The medicine is coated for easier swallowing. (_ _ _ _)



B. Answer the questions.

1. What is the form of this medicine?
2. What is the active ingredient in this product?
3. How should we use this medicine? Can we take it orally?
4. What is the main purpose of this rubbing alcohol?
5. How many fluid ounces (fl. oz.) are in the bottle?
6. Why must this product be kept away from fire or flame?
7. According to the label, how often should you apply this product on the affected area?
8. A person gets a small cut on their hand. How should they use this product according to the directions?
9. What should you do before using a bandage after applying this rubbing alcohol?
10. How should this product be stored?
11. If the safety foil under the cap is broken or missing, what should you do?