

DENOMINADOR COMÚN

$$M(2) = \{ _, _, _, _ \dots \}$$

$$M(8) = \{ _, _, _, _ \dots \}$$

2	8	:

mcm:

$$M(3) = \{ _, _, _, _ \dots \}$$

$$M(15) = \{ _, _, _, _ \dots \}$$

3	15	:

mcm:

$M(2) = \{ _, _, _, _ \dots \}$

$M(5) = \{ _, _, _, _ \dots \}$

2	5	:

mcm:

$M(4) = \{ _, _, _, _ \dots \}$

$M(10) = \{ _, _, _, _ \dots \}$

4	10	:

mcm:

FRACCIONES EQUIVALENTES POR MEDIO DE LA AMPLIFICACIÓN

$$\frac{1}{2} \cdot \frac{2}{2} = \frac{\quad}{\quad}$$

$$\frac{1}{2} \cdot \frac{3}{3} = \frac{\quad}{\quad}$$

$$\frac{1}{2} \cdot \frac{4}{4} = \frac{\quad}{\quad}$$

$$\frac{1}{2} \cdot \frac{5}{5} = \frac{\quad}{\quad}$$

$\frac{1}{2} \rightarrow$ denominador 6 $\rightarrow$

$\frac{1}{4} \rightarrow$ denominador 8 $\rightarrow$

$\frac{4}{5} \rightarrow$ denominador 20 $\rightarrow$

$\frac{2}{3} \rightarrow$ denominador 9 $\rightarrow$

AMPLIFICACIÓN PARA
OBTENER UN DENOMINADOR