

Time: 1Hour

Marks :40

Four alternatives are given for each of the incomplete statement or questions. Choose the correct answer.

- 1) The lines representing $2x + 3y - 9 = 0$ & $4x + 6y - 18 = 0$ are
A) Intersecting lines B) Perpendicular lines C) Parallel lines D) Coincident lines
- 2) 10th term of an A.P. 5,9,13,..... is
A) 36 B) 31 C) 41 D) 21
- 3) Roots of the equation $ax^2 + bx + c = 0$ are
A) $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ B) $x = \frac{b \pm \sqrt{b^2 - 4ac}}{2a}$ C) $x = \frac{-b \pm \sqrt{b^2 + 4c}}{2a}$ D) $x = \frac{-b - \sqrt{b^2 - 4ac}}{2}$
- 4) Value of $\sin 30^\circ + \cos 60^\circ$ is
A) $\frac{1}{2}$ B) $\frac{3}{2}$ C) $\frac{1}{4}$ D) 1
- 5) Distance between the origin and a point (3,4) is
A) 5 B) 6 C) -5 D) -4
- 6) A line passing through 2 points on the circumference of a circle is
A) Chord B) Secant C) Tangent D) Radius
- 7) Total Surface Area of a hemisphere is
A) πr^2 B) $4 \pi r^2$ C) $\frac{4}{3} \pi r^2$ D) $3 \pi r^2$
- 8) If area of the circular base is 154 cm^2 , height = 10cm, then the volume of the cylinder is
A) 15.40 cm^3 B) 15400 cm^3 C) 1.540 cm^3 D) 1540 cm^3
- 9) If $a_1x + b_1y + c_1 = 0$ and $a_2x + b_2y + c_2 = 0$ lines are inconsistent lines, then the ratio of their coefficients is
A] $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$ B] $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$ C] $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$ D] $\frac{a_1}{a_2} = \frac{b_1}{b_2}$
- 10) Sum of first 10 natural numbers is
A) 55 B) 50 C) 40 D) 10
- 11) If a quadratic equation has two equal roots, then the discriminant is
A) $b^2 - 4ac < 0$ B) $b^2 - 4ac > 0$ C) $b^2 - 4ac = 0$ D) $b^2 - 4ac \geq 0$
- 12) Value of $\sin^2\theta + \cos^2\theta$ is
A) 0 B) 1 C) 2 D) 3
- 13) The coordinates of a point P(x,y) which divides the line segment joining the points A(x_1, y_1) and B(x_2, y_2) in the ratio $m_1 : m_2$ is

- A] $\left(\frac{m_1 x_2 + m_2 x_1}{m_1 + m_2}, \frac{m_1 y_2 + m_2 y_1}{m_1 + m_2} \right)$ B] $\left(\frac{m_1 x_2 - m_2 x_1}{m_1 + m_2}, \frac{m_1 y_2 - m_2 y_1}{m_1 + m_2} \right)$
C] $\left(\frac{m_1 x_2 + m_2 x_1}{m_1 - m_2}, \frac{m_1 y_2 + m_2 y_1}{m_1 - m_2} \right)$ D] $\left(\frac{m_1 x_1 + m_2 x_2}{m_1 + m_2}, \frac{m_2 y_2 + m_2 y_2}{m_1 + m_2} \right)$

14) Tangents drawn at the ends of a diameter are mutually

- A) Perpendicular B) Parallel C) Intersecting D) Coinciding

15) Curved surface area of Cone is

- A) πrh B) $4\pi r(r+l)$ C) πrl D) $\pi r^2 l$

16) Corresponding sides of two similar triangles are in the ratio 1: 4. Areas of these triangles are in the ratio

- A) 1:2 B) 1:16 C) 1:4 D) 16:1

17) If $4x+py+8=0$ and $4x+4y+2=0$, represent parallel lines, the value of p is

- A) 1 b) 2 C) 4 d) 8


18) The discriminant of a quadratic equation $x^2 -5x + 3 = 0$ is

- A) 25 B) 20 C) 13 D) -13

19) Value of $\frac{\sin 28}{\cos 62}$ is

- A) 0 B) 1 C) 2 D) 28

20) In $\triangle ABC$, $AB \perp BC$, $\angle ACB = 30^\circ$, $AB = 15\sqrt{3}m$, value of BC is



- A) $15\sqrt{3}m$ B) 10m C) 45m D) 50m

21) Midpoint of the line joining the points (6,2) and (4,4) is

- A) (5,2) B) (3,5) C) (2,5) D) (5,3)

22) If mean and median are 3 and 4 respectively, then the mode is

- A) 12 B) 6 C) 4 D) 3

23) In a right triangle square of the _____ is equal to the sum of the squares of other two sides.

- A) Perpendicular B) sum C) Hypotenuse D) Opposite

24) Formula to find the curved surface area of a Cylinder is

- A) πrh B) $4\pi r(r+l)$ C) $2\pi r(r+h)$ D) $2\pi rh$

25) 4 pens and 5 pencils together cost Rs.25. Represent this in the form of linear equation in two variables.

- A) $4x + 5y = 25$ B) $x + y = 25$ C) $x - y = 25$ D) $4x - 5y = 25$

26) Roots of $x^2 - 25 = 0$ are

- A) 25 and - 25 B) 5 and 0 C) -5 and 0 D) 5 and -5

- 27) The value of $\sin 90^\circ + \tan 45^\circ$.
- A) 0 B) 1 C) 2 D) 3
- 28) If $\sec \theta = \frac{1}{\sqrt{5}}$, then $\cos \theta$ is
- A) 0 B) 1 C) 5 D) $\sqrt{5}$
- 29) The distance of the co-ordinate P(3,4) from the x-axis is
- A) 3 B) 4 C) 5 D) 1
- 30) Median of the scores 15,17,19,21,23,25 is
- A) 19 B) 20 C) 21 D) 6
- 31) If a line is drawn parallel to one side of a triangle to intersect the other two sides in distinct points, then the other two sides are divided in the
- A) same point B) same ratio C) same distance D) same length
- 32) If the volume of a Cylinder is 300cm^3 , then volume of the cone of same base and height is
- A) 300cm^3 B) 200cm^3 C) 100cm^3 D) 50cm^3
- 33) If $a_n = 3n - 2$, then the value of a_4 is,
- A) 10 B) 12 C) 8 D) 6
- 34) Which one of the following are similar figures?
- A) Rectangles B) Rhombuses C) Trapeziums D) Circles
- 35) Maximum number of tangents that can be drawn to a circle from an external point is
- A) Infinite B) 1 C) 2 D) 0
- 36) In a right triangle ABC, if $\angle B = 90^\circ$, $AB = 5\text{cm}$, $BC = 12\text{cm}$, the length of AC is
- A) 10cm B) 12cm C) 13cm D) 17cm
- 37) Angle formed by the line of sight with the horizontal when the point being viewed is above the horizontal level is
- A) angle of elevation B) angle of depression C) equal angle D) Right angle
- 38) Mean of 5,6,7,8,9,10,11 is
- A) 5 B) 8 C) 11 D) 45
- 39) We plot lower limits on the x-axis and the corresponding cumulative frequencies on the y axis in
- A) Histogram B) Pie chart C) Less than Ogive D) More than Ogive
- 40) If a Sphere is melted and recast in to the shape of a cylinder, the thing which remains same is,
- A) Length B) Radius C) Area D) Volume