

**Pre-Assessment****Name:****Grade & Section:****Choose the best answer from the options that follow each question.**

1. A 15.0 mL volume of 0.0100 M  $\text{Pb}(\text{NO}_3)_2$  is added to 15.0 mL of 0.0100 M  $\text{NaI}$ . A precipitate is formed, and equilibrium is established. The  $K_{sp}$  equilibrium expression for the dissolution of  $\text{PbI}_2$  is

- $[\text{Pb}^{2+}][\text{I}^-]^2$ .
- $[\text{Pb}][\text{I}^2]$ .
- $[\text{Pb}^{2+}][2\text{I}^-]^2$ .
- $2[\text{Pb}^{2+}][\text{I}^-]$ .

2. A 15.0 mL volume of 0.0100 M  $\text{Pb}(\text{NO}_3)_2$  is added to 15.0 mL of 0.0100 M  $\text{NaI}$ . A precipitate is formed, and equilibrium is established. What is the ion product for the reaction that produces  $\text{PbI}_2$ ?

- $1.00 \times 10^{-8}$
- $1.25 \times 10^{-7}$
- $1.00 \times 10^{-6}$
- $2.5 \times 10^{-5}$

3. What is the equilibrium expression for the following equation?

$$\text{Fe(OH)}_3(aq) \rightleftharpoons \text{Fe}^{3+}(aq) + 3\text{OH}^-(aq)$$

- $$\frac{[\text{Fe}^{3+}][\text{OH}^-]}{[\text{Fe(OH)}_3]}$$
- $$\frac{[\text{Fe}^{3+}][\text{OH}^-]^3}{[\text{Fe(OH)}_3]}$$
- $$\frac{[\text{Fe(OH)}_3]}{[\text{Fe}^{3+}][\text{OH}^-]^3}$$
- $$\frac{[\text{Fe(OH)}_3]}{[\text{Fe}^{3+}][\text{OH}^-]}$$

4. In a bottle of unopened cola, the  $\text{CO}_2$  gas dissolved in the liquid is in equilibrium with the  $\text{CO}_2$  gas above the liquid. The dissolved gas reacts with water molecules in the cola to form carbonic acid, which also dissociates into carbon dioxide and water. Which chemical equation(s) best describe this equilibrium system?

- $\text{CO}_2(g) \rightleftharpoons \text{CO}_2(l)$
- $\text{CO}_2(g) \rightleftharpoons \text{CO}_2(aq)$  and  $\text{CO}_2(l) + \text{H}_2\text{O}(l) \rightleftharpoons \text{H}_2\text{CO}_3(aq)$
- $\text{CO}_2(g) \rightleftharpoons \text{CO}_2(aq)$
- $\text{CO}_2(g) \rightleftharpoons \text{CO}_2(aq)$  and  $\text{CO}_2(aq) + \text{H}_2\text{O}(l) \rightleftharpoons \text{H}_2\text{CO}_3(aq)$

5. The Haber process,  $\text{N}_2(g) + 3\text{H}_2(g) \rightleftharpoons 2\text{NH}_3(g) + 92 \text{ kJ}$ , is operated at temperatures of about  $500^\circ\text{C}$  because


- a higher temperature would favor the products.
- $K$  is a maximum at  $500^\circ\text{C}$ .
- the forward reaction rate is too slow at lower temperatures.
- $K$  is equal to 1 at  $500^\circ\text{C}$ .

6. Which of the following would not affect the aqueous equilibrium reaction represented by the equation below?



- a. the addition of  $\text{NaNO}_3(s)$
- b. an increase in  $\text{H}^+$  concentration
- c. the addition of  $\text{NaNO}_2(s)$
- d. a decrease in  $\text{NO}_2^-$  concentration

7. Consider the equilibrium system represented by the equation below.



If the hydroxide ions were removed, how would the color change?

If the hydroxide ions

- a. to darker yellow
- b. to lighter orange
- c. to darker orange
- d. not at all

8. The  $\text{Fe}^{3+}$  ion present in acid mine drainage is colorless and surrounded by water molecules. If phenanthroline (Phe) is added, the solution turns orange as a colored complex is formed according to the equation  $\text{Fe}^{3+} \cdot \text{H}_2\text{O} + \text{Phe} \rightleftharpoons \text{Fe}^{3+} \cdot \text{Phe} + \text{H}_2\text{O}$ . The color is commonly used as an indicator of the  $\text{Fe}^{3+}$  ion concentration. What would an increase in color from light to dark orange indicate in this system?

- an applied stress that shifted the equilibrium to favor the reverse reaction
- an applied stress that shifted the equilibrium to favor the forward reaction
- an increase in the rate of the reverse reaction
- addition of water to the system

9. The solubility product constant expression includes

- a concentration of zero for undissolved salt.
- the concentrations of dissolved and undissolved salt.
- an exponent corresponding to the charge on each ion.
- the concentrations of dissociated ions.

10. In a reaction that goes nearly to completion,

- the re-formation of reactants is much slower than the formation of the products.
- the rate of the reverse reaction is faster than the rate of the forward reaction.
- the re-formation of reactants is much faster than the rate of formation of the products.
- the forward reaction rate increases.

11. Which of the following  $K_{eq}$  expressions is correct for the formation of ammonia,  $\text{NH}_3$ , from its elements, as represented by the equation  
$$3\text{H}_2 + \text{N}_2 \rightleftharpoons 2\text{NH}_3 + 92 \text{ kJ}$$

- $$\frac{[\text{NH}_3]^2}{[\text{H}_2]^3[\text{N}_2]}$$
- $$\frac{[\text{H}_2]^3[\text{N}_2]}{[\text{NH}_3]^2}$$
- $$\frac{[\text{NH}_3]}{[\text{H}_2][\text{N}_2]}$$
- $$\frac{[\text{H}_2][\text{N}_2]}{[\text{NH}_3]}$$

12. When you calculate the  $K_{sp}$  of calcium fluoride,  $\text{CaF}_2$ , if the concentration of the  $\text{F}^-$  ion is  $4.2 \times 10^{-4} \text{ M}$ , then the concentration of the  $\text{Ca}^{2+}$  ion is

- $2.1 \times 10^{-4} \text{ M}$ .
- $8.4 \times 10^{-4} \text{ M}$ .
- $4.2 \times 10^{-4} \text{ M}$ .
- $3.7 \times 10^{-11} \text{ M}$ .

13. Which of the following salts is *least* soluble?

- $\text{Ag}_2\text{CO}_3$   $K_{sp} = 8.4 \times 10^{-12}$
- $\text{Ag}_2\text{CrO}_4$   $K_{sp} = 1.1 \times 10^{-12}$
- $\text{AgI}$   $K_{sp} = 1.5 \times 10^{-16}$
- $\text{AgBr}$   $K_{sp} = 5.4 \times 10^{-13}$

14. Consider the reaction represented by the equation  $2\text{NO}(g) + \text{O}_2(g) \rightleftharpoons 2\text{NO}_2(g)$ . If the volume of the reaction chamber is decreased, then the

- forward reaction will be favored.
- reverse reaction will be favored.
- initial pressure of  $\text{NO}_2$  will decrease.
- initial pressure of  $\text{O}_2$  will remain constant.

15. Consider the reaction represented by the equation  $\text{Ag}_2\text{SO}_4(aq) \rightleftharpoons 2\text{Ag}^+(aq) + \text{SO}_4^{2-}(aq)$ . You can shift the equilibrium to favor the reverse reaction by adding

- $\text{CaCl}_2$ .
- $\text{AgNO}_3$ .
- $\text{Na}_2\text{SO}_4$ .
- Both (b) and (c)

16. A chemical reaction that is at equilibrium always has

- a high  $K_{eq}$  value.
- a forward reaction rate that equals the reverse reaction rate.
- equal concentrations of reactants and products.
- Both (a) and (b)

\_\_\_\_ 17. Which of these is true about the chemical equation below?  
reactants  $\rightleftharpoons$  products

- The concentration of the products is greater than the concentration of the reactants.
- $K_{eq}$  of the forward reaction has a low value.
- The reaction favors the products.
- $K_{eq}$  is equal to 1.

\_\_\_\_ 18. In which of the following reactions, described by the equations below, will an increase in pressure have no effect on the equilibrium of the system?

- $2\text{NO}_2(g) \rightleftharpoons \text{N}_2\text{O}_4(g)$
- $2\text{NOCl}(g) \rightleftharpoons 2\text{NO}(g) + \text{Cl}_2(g)$
- $\text{H}_2\text{O}(g) + \text{CO}(g) \rightleftharpoons \text{H}_2(g) + \text{CO}_2(g)$
- $\text{N}_2(g) + 3\text{H}_2(g) \rightleftharpoons 2\text{NH}_3(g)$

\_\_\_\_ 19. The common-ion effect promotes

- dissolution.
- precipitation.
- boiling.
- ionization.

\_\_\_\_ 20. What is the acid ionization expression for the equation  $\text{HA}(aq) + \text{H}_2\text{O}(l) \rightleftharpoons \text{H}_3\text{O}^+(aq) + \text{A}^-(aq)$ ?

- $$\frac{[\text{A}^-][\text{H}_3\text{O}^+]}{[\text{HA}][\text{H}_2\text{O}]}$$
- $$\frac{[\text{A}^-]}{[\text{HA}][\text{H}_2\text{O}]}$$
- $$\frac{[\text{A}^-][\text{H}_3\text{O}^+]}{[\text{HA}]}$$
- $$[\text{A}^-][\text{H}_3\text{O}^+][\text{HA}][\text{H}_2\text{O}]$$

\_\_\_\_ 21. The reaction represented by the equation  $\text{BH}^+(aq) + \text{H}_2\text{O}(l) \rightleftharpoons \text{H}_3\text{O}^+(aq) + \text{B}(aq)$  is an example of a(n)

- cation hydrolysis reaction.
- anion hydrolysis reaction.
- conjugated reaction.
- pH reaction.

\_\_\_\_ 22. What is the value of  $K_w$ ?

- $1 \times 10^{-14}$
- $1 \times 10^{14}$
- $1 \times 10^7$
- $14 \times 10^{-14}$

\_\_\_\_ 23. An example of a good buffer solution is one that contains

- HCl and NaCl.
- $\text{HNO}_2$  and NaCl.
- $\text{HNO}_2$  and  $\text{NaNO}_2$ .
- $\text{CH}_3\text{COOH}$  and NaCl.

\_\_\_\_ 24. When  $\text{H}_3\text{O}^+$  ions are added to an aqueous solution of acetic acid,

- the reaction forms more  $\text{CH}_3\text{COO}^-$ .
- the reaction forms more water.
- the reaction forms more  $\text{CH}_3\text{COOH}$ .
- nothing happens.

\_\_\_\_ 25. Which of the following is a conjugate acid-base pair in the reaction represented by the equation below?

$$\text{H}_2\text{PO}_4^- + \text{H}_2\text{O} \rightleftharpoons \text{H}_3\text{PO}_4 + \text{OH}^-$$

- $\text{H}_2\text{PO}_4^-$  and  $\text{H}_2\text{O}$
- $\text{H}_2\text{PO}_4^-$  and  $\text{OH}^-$
- $\text{H}_2\text{PO}_4^-$  and  $\text{H}_3\text{PO}_4$
- None of the above

