

**Drag the correct number to the correct question on the following prescription label.**

| <b>Drug Facts</b>   |  |
|---|--|
| <b>Active ingredient (in each tablet)</b><br>Chlorpheniramine maleate 2 mg.....   | <b>Purpose</b><br>Antihistamine  |
| <b>Uses</b> temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ sneezing ■ runny nose ■ itchy, watery eyes ■ itchy throat  |  |
| <b>Warnings</b><br>Ask a doctor before use if you have<br>■ glaucoma ■ a breathing problem such as emphysema or chronic bronchitis<br>■ trouble urinating due to an enlarged prostate gland<br>Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives |  |
| <b>When using this product</b><br>■ drowsiness may occur ■ avoid alcoholic drinks<br>■ alcohol, sedatives, and tranquilizers may increase drowsiness<br>■ be careful when driving a motor vehicle or operating machinery<br>■ excitability may occur, especially in children      |  |
| If pregnant or breast-feeding, ask a health professional before use.<br>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.   |  |
| <b>Directions</b>   |  |
| adults and children 12 years and over   | take 2 tablets every 4 to 6 hours;<br>not more than 12 tablets in 24 hours |
| children 6 years to under 12 years  | take 1 tablet every 4 to 6 hours;<br>not more than 6 tablets in 24 hours   |
| children under 6 years  | ask a doctor   |

|   |
|---|
| <b>Drug Facts (continued)</b>   |
| <b>Other information</b> ■ store at 20-25°C (68-77°F) ■ protect from excessive moisture                                       |
| <b>Inactive ingredients</b> D&C yellow no. 10, lactose, magnesium stearate, microcrystalline cellulose, pregelatinized starch |

1. What is the purpose of this drug?

Drag Answer Here

2. Children Under 6 years old should never take this drug.

TRUE

FALSE

## Drag the correct number to the correct question on the following prescription label.

3. An adult or child who is 12 years or older should take this medication how often?

Drag Answer Here

4. A pregnant women should?

Drag Answer Here

5. \_\_\_\_\_ may occur, especially in small children.

Drag Answer Here

6. If you have the following you should ask a doctor before using this medication. CHECK all that apply.

Glaucoma

Enlarged  
Prostrate

Bronchitis

Emphysema

Sneezing

7. Avoid \_\_\_\_\_ drinks.

Drag Answer Here

8. Children under the age of 12 and at least 6 years old should take this medication.....

Drag Answer Here

9. You should go ahead and take this medication if you are currently taking a sedative.

TRUE

FALSE

10. What is the active ingredient in this medication

Drag Answer Here